



Selectivity filter instability dominates the low intrinsic activity of the TWIK-1 K2P K⁺ channel

Received for publication, August 14, 2019, and in revised form, December 4, 2019. Published, Papers in Press, December 5, 2019, DOI 10.1074/jbc.RA119.010612

Ehsan Nematian-Ardestani[‡], Firdaus Abd-Wahab^{‡1}, Franck C. Chatelain[§], Han Sun[¶],  Marcus Schewe^{||}, Thomas Baukrowitz^{||}, and  Stephen J. Tucker^{‡2}

From the [‡]Clarendon Laboratory, Department of Physics, University of Oxford, Oxford OX1 3PU, United Kingdom, the [§]Université Côte d'Azur, Centre National de la Recherche Scientifique, Institut de Pharmacologie Moléculaire et Cellulaire, Labex ICST, 06560 Valbonne, France, the [¶]Department of Structural Biology, Leibniz-Forschungsinstitut für Molekulare Pharmakologie, 13125 Berlin, Germany, and the ^{||}Institute of Physiology, Christian-Albrechts University of Kiel, 24118 Kiel, Germany

Edited by Mike Shipston

Two-pore domain K⁺ (K2P) channels have many important physiological functions. However, the functional properties of the TWIK-1 (K2P1.1/*KCNK1*) K2P channel remain poorly characterized because heterologous expression of this ion channel yields only very low levels of functional activity. Several underlying reasons have been proposed, including TWIK-1 retention in intracellular organelles, inhibition by posttranslational sumoylation, a hydrophobic barrier within the pore, and a low open probability of the selectivity filter (SF) gate. By evaluating these potential mechanisms, we found that the latter dominates the low intrinsic functional activity of TWIK-1. Investigating this further, we observed that the low activity of the SF gate appears to arise from the inefficiency of K⁺ in stabilizing an active (*i.e.* conductive) SF conformation. In contrast, other permeant ion species, such as Rb⁺, NH₄⁺, and Cs⁺, strongly promoted a pH-dependent activated conformation. Furthermore, many K2P channels are activated by membrane depolarization via an SF-mediated gating mechanism, but we found here that only very strong nonphysiological depolarization produces voltage-dependent activation of heterologously expressed TWIK-1. Remarkably, we also observed that TWIK-1 Rb⁺ currents are potently inhibited by intracellular K⁺ (IC₅₀ = 2.8 mM). We conclude that TWIK-1 displays unique SF gating properties among the family of K2P channels. In particular, the apparent instability of the conductive conformation of the TWIK-1 SF in the presence of K⁺ appears to dominate the low levels of intrinsic functional activity observed when the channel is expressed at the cell surface.

K2P³ K⁺ channels represent a structurally unique family of channels involved in diverse physiological functions such as cell volume regulation, apoptosis, vasodilation, central chemosen-

sitivity, neuronal excitability, and perception of pain. They also act as a major target for volatile anesthetics and represent attractive therapeutic targets for the treatment of a wide variety of cardiovascular and neurological disorders (1–3).

The first mammalian member of this family to be identified was the TWIK-1 (K2P1/*KCNK1*) channel (4). This channel has subsequently been implicated in a number of important physiological roles in the heart and brain (5–7). However, understanding the biophysical, pharmacological, and functional properties of TWIK-1 has proven difficult because of the extremely low levels of functional activity observed when the WT channel is recombinantly expressed in heterologous systems such as *Xenopus* oocytes or cultured mammalian cells (*e.g.* HEK293) that are typically used for electrophysiological recordings of ion channels.

The precise reasons underlying these low levels of functional activity remain unclear. Originally, it was suggested that post-translational modification (sumoylation) of a lysine residue (Lys²⁷⁴) within the proximal C terminus was responsible for its apparent lack of functional activity (8). The structure of TWIK-1 revealed Lys²⁷⁴ to be located within an amphipathic interfacial “C-helix” structure kinked parallel to the membrane, close to the intracellular entrance to the pore (9). However, the mechanisms underlying the increased activity seen with mutations at Lys²⁷⁴ within the C-helix remain controversial (10).

A subsequent study of WT TWIK-1 channels revealed that they are actively and constitutively internalized from the cell surface and that mutation of an endocytic motif within the intracellular C-terminal domain can reduce this internalization to increase channel levels within the plasma membrane (11). However, even when engineered to reside on the cell surface in greater numbers, these “WT” TWIK-1 channels still exhibit very low levels of functional activity compared with many other members of the K2P channel family.

Exploiting this approach to stabilize TWIK-1 within the plasma membrane, it was also found that larger TWIK-1 currents can be produced by mutation of hydrophobic residues that form a “hydrophobic cuff” within the inner pore of the channel (*e.g.* Leu¹⁴⁶). This cuff creates a hydrophobic barrier that restricts hydration of the inner cavity below the selectivity filter (12–15). However, the precise role of this barrier also remains unclear because, in addition to its effect on pore hydration, mutation of residues that form this barrier may also indi-

This work was supported by Biotechnology and Biological Sciences Research Council Grant BB/N000145/1 (to E. N.-A. and S. J. T.), Deutsche Forschungsgemeinschaft Research Group FOR 2518 Dynlon (to H. S., M. S., and T. B.), and CNRS. The authors declare that they have no conflicts of interest with the contents of this article.

This article contains Figs. S1–S3.

¹ Supported by a King's scholarship from the Malaysian government.

² To whom correspondence should be addressed. E-mail: stephen.tucker@physics.ox.ac.uk.

³ The abbreviations used are: K2P, two-pore domain K⁺; SF, selectivity filter; TPA, tetrapentylammonium; MD, molecular dynamics.




Figure 1. Role of charged residues within the C-helix. *A*, -fold-increase in activity of TWIK-1 mutations in the C-helix compared with a mutation within the hydrophobic cuff (L146N). Maximum whole-cell currents were recorded at 0 mV from channels expressed in *Xenopus* oocytes. All mutations were introduced into the TWIK-1* construct, which exhibits greater stability within the plasma membrane. The dotted red line indicates the level of current recorded for WT TWIK-1* ($0.8 \pm 0.1 \mu\text{A}$ at 0 mV, $n \geq 18$; see also Fig. S1). *B*, the three activatory mutations within the C-helix are positively charged and point toward the inner mouth of the pore. Mutation of other charged residues that point away does not increase currents.

rectly affect other aspects of K₂P channel gating (16). Overall, detailed functional characterization of nonmutant TWIK-1 channels remains challenging.

Another unusual structural feature of the TWIK-1 channel is its highly asymmetric and noncanonical selectivity filter (SF). By nature of their dimeric architecture, all K₂P channels exhibit a two-fold symmetry and have a SF formed from nonidentical sequences, but for TWIK-1, there are also major variations in the canonical TXGY/FG sequence, with TTGYG in the first P-loop and TIGLG in the second. Furthermore, the channel has been shown to exhibit dynamic ion selectivity with reduced K⁺ > Na⁺ selectivity that can be modulated by mutations either within or close to the first P-loop (15, 17, 18). Interestingly, this dynamic ion selectivity is thought to be relevant for regulation of cardiac excitability under conditions of hypokalemia (5, 7).

In most other K₂P channels, the SF also represents the main site for channel gating and the origin of their voltage sensitivity (19–22). In a previous study, we demonstrated that this voltage-dependent gating originates from the movement of ions into the high electric field of the SF to generate a one-way “check valve” within the filter, where the outward movement of K⁺ induces filter opening and inward movement promotes channel closure. The gating process is also highly dependent on the nature of the permeant ion (23). However, at that time, the relevance of this filter gating mechanism to TWIK-1 was not examined because of its apparent linear current–voltage behavior within the physiological voltage range. Nevertheless, it has been shown previously that macroscopic TWIK-1 currents could be recorded when Rb⁺ was used as the permeant ion (12, 17), suggesting the presence of some type of filter-gating mechanism similar to that seen in other K₂P channels.

In this study, we therefore sought to understand the extent to which these different possible mechanisms contribute to the low levels of functional activity associated with TWIK-1. We found that TWIK-1 does indeed possess a filter gate that displays characteristics markedly different from most other K₂P channels and that this appears to be the principal reason why TWIK-1 produces very low levels of channel activity when expressed heterologously.

Results

Role of the C-helix

To begin to evaluate the effect of these different mechanisms, we first examined the role of the C-helix. A lysine residue within the C-helix (Lys²⁷⁴) has been proposed to be a site for posttranslational sumoylation that silences TWIK-1 activity (8). However, the C-helix also contains a number of other charged residues. We therefore conducted a scanning mutagenesis of the C-helix and compared the levels of activity produced when these mutant channels were expressed in *Xenopus* oocytes.

Constitutive endocytosis has been reported previously to reduce the levels of TWIK-1 within the plasma membrane (11, 15). Therefore, to permit measurement of larger currents, all mutations were created in a background construct where the di-isoleucine retention motif was mutated (I293A/I294A). This site is located within the cytoplasmic domain and is consistent with many previous studies of TWIK-1; this trafficking mutant (hereafter referred to as TWIK-1*) was used throughout this study to measure channel activity.

As reported previously (12), we found that mutation of Lys²⁷⁴ in TWIK-1* caused an increase in whole-cell currents. However, although we found that most other mutations in the C-helix had little effect, mutation of two other charged residues (Arg²⁷⁷ and Lys²⁷⁸) also increased the level of whole-cell current activity (Fig. 1A and Fig. S1). Interestingly, when mapped onto the structure of the C-helix, these three charged residues were observed to face toward the entrance of the inner cavity. By contrast, mutation of adjacent charged residues (e.g. Glu²⁷² and Lys²⁷⁵) that face away from the pore had no such effect (Fig. 1, A and B).

The precise mechanisms responsible for the increased activity of these pore-facing mutations remains unclear and likely involves multiple factors. However, the fact that these three pore-facing mutants were all able to increase macroscopic current levels to the same levels as mutation of Lys²⁷⁴ strongly suggests that the role of Lys²⁷⁴ in “silencing” this channel may not be as unique as thought previously.

Importantly, and consistent with previous reports, the increase in currents seen with these C-helix mutants is also relatively small compared with the much bigger effect of mutations within the hydrophobic cuff of the inner pore, e.g. the L146N mutation in M2 (Fig. 1A). Furthermore, we have also

Unusual filter gating properties of TWIK-1




Figure 2. Functional characterization of TWIK-1* channels in excised patches. *A*, example of K⁺ currents recorded from TWIK-1* in giant excised patches in response to voltage steps from -100 mV to $+200$ mV. Note the supraphysiological voltages (>100 mV) required to observe voltage-dependent activation. At $+140$ mV, TWIK-1* currents averaged 0.49 ± 0.1 nA compared with 0.01 ± 0.005 nA for uninjected controls ($n \geq 6$). *B*, comparison of the currents recorded in giant excised patches from oocytes expressing WT TWIK-1 or the TWIK-1* trafficking mutant I293A/I294A. Currents were recorded in response to a voltage ramp from -70 mV to $+70$ mV. Patches have 120 mM Rb⁺ on the inside with 120 mM K⁺ in the pipette solution. *C*, *left panel*, activation of TWIK-1* currents when K⁺ is replaced with Rb⁺ on the extracellular side. Currents were recorded in excised patches from oocytes in response to a series of voltage steps from $+120$ to -120 mV with 120 mM Rb⁺ in the pipette and 120 mM intracellular K⁺. *Right panel*, a current-voltage plot from this example recording. *D*, intracellular pH sensitivity of TWIK-1* Rb⁺ currents. *E* and *F*, normalized TWIK-1* Rb⁺ currents plotted against intracellular pH (*E*) and plotted against TPA concentration (*F*).

shown that both WT TWIK-1* and L146N mutant channels produce even larger macroscopic currents when Rb⁺ is used as the permeant ion (12). This indicates that the low levels of TWIK-1* activity typically observed when the channel is expressed can only partly be accounted for by the C-helix, and the intrinsic biophysical properties of the SF are likely to play a more dominant role in this process.

Functional properties of TWIK-1 currents using Rb⁺ as the permeant ion

Our previous studies identified an ion flux-gating mechanism within the SF of many different subfamilies of K2P channels (23), but its contribution to TWIK-1 gating remains unclear. We therefore next examined how the SF gate contributes to the unusual functional properties of TWIK-1.

When K⁺ is used as the permeant ion with either WT TWIK-1 or TWIK-1*, relatively little current can normally be recorded above background levels within a physiological voltage range (*i.e.* -100 to $+100$ mV) (4, 23). However, Fig. 2A shows that, in excised patches, strong depolarization in excess of $+100$ mV begins to elicit measurable K⁺ currents compared with uninjected controls. These currents display similar time-

and voltage-dependent activation properties observed in other K2P channels, such as TREK-1 and TRAAK. However, unlike these other K2P channels, the threshold for this activation is shifted to >100 mV, highlighting the unusual flux-gating properties of the SF gate in TWIK-1*.

When Rb⁺ is used as the permeant ion in TREK and TRAAK channels, it activates the SF ion flux gating mechanism within the SF by stabilization of its active configuration (23, 24). Consistent with this, we also observed that Rb⁺ is capable of producing large currents in inside-out macropatches excised from *Xenopus* oocytes expressing WT TWIK-1*. These currents were also activated at less depolarized potentials than required to generate K⁺ currents (Fig. 2B). This suggests that K⁺ is much less efficient than Rb⁺ in stabilizing the SF in an actively conductive configuration. Fig. 2B also highlights the relative importance of the endocytotic retention motif because, when the retention motif is present (*i.e.* WT TWIK-1), then the currents are markedly smaller, even with Rb⁺ as the permeant ion. The ability to record robust Rb⁺ currents from WT TWIK-1* therefore provides an opportunity for a more detailed biophysical investigation of the intrinsic gating properties of this unusual K2P channel.




Figure 3. Role of different permeant ions. *A*, examples of currents recorded in a single giant excised inside-out patch from an oocyte expressing TWIK-1* in response to a series of voltage steps from -120 to $+140$ mV. The indicated ions (120 mM each) were exchanged on the intracellular side. Traces are ordered based on the potency of activation by each ion. *B*, current voltage plots for each of these different ions. *C*, the rate of activation by Rb⁺ appears to be independent of the applied voltage. *D*, comparison of TWIK-1* Rb⁺ activation at room temperature and at 3°C . Despite the obvious decrease in the rate of activation, no tail currents are observed at 3°C (see also Fig. S2B).

In the excised macropatch recordings described above, Rb⁺ is applied intracellularly, and its activatory effect is consistent with the intracellular effects of Rb⁺ seen for many other K2P channels. However, consistent with previous reports (17), we also observed that this activatory effect occurs extracellularly, *i.e.* when Rb⁺ is included in the pipette solution (Fig. 2C). This is in contrast to many other K2P channels (*e.g.* TREK-1), where Rb⁺ activation only works from the intracellular side (23). This provides further evidence that the TWIK-1 SF gating mechanism is different from most other members of the K2P channel family.

TREK/TRAAK channel activity can also be regulated by intracellular H⁺, and so we examined whether intracellular pH also regulates TWIK-1* activity. Fig. 2, D and E, shows that the robust Rb⁺ currents recorded from TWIK-1* in excised patches are also sensitive to inhibition by intracellular H⁺ (pH_{0.5}, ~ 7.8). Thus, to optimize currents, all experiments were subsequently performed using an intracellular pH of 8.0. Furthermore, to distinguish TWIK-1* currents from leak, we quantified the effects of intracellular tetrapentylammonium (TPA) as a channel blocker (Fig. 2F). We found that intracellular TPA blocked TWIK-1* Rb⁺ currents with an even higher affinity (IC₅₀ = 3.3 ± 0.1 μM) than typically observed for most other K2P channels (21, 25). These results therefore established an experimental framework for the measurement of robust TWIK-1 channel activity.

Stabilization of the SF gate by different permeant ions

Several other monovalent cations have also been shown to activate the ion flux gating mechanism in other K2P channels, with Rb⁺ and NH₄⁺ typically being the most potent and Na⁺ the least (23). Fig. 3, A–C, and Fig. S2 show a similar pattern of activation for WT TWIK-1*. But, in marked contrast to other K2Ps where Cs⁺ acts as a permeant blocker, Cs⁺ appears to activate the SF gate in TWIK-1* more effec-

tively than K⁺. Furthermore, despite the levels of activation induced by intracellular NH₄⁺ and Rb⁺, no tail currents could be observed upon repolarization (Fig. 3, A and D). This is in marked contrast to many other K2P channels where, under similar conditions, large tail currents can be observed upon repolarization (23).

The activation of TWIK-1* by extracellular Rb⁺ (Fig. 2C) rules out the possibility that the channel cannot conduct extracellular ions into the cell, but to further examine the apparent absence of tail currents, we repeated these measurements at 3°C to slow down channel gating kinetics. As a control for this effect, we used a mutant version of TREK-2 (26) where the normally large tail currents have become so fast as to appear absent at 23°C , but reducing the temperature slowed the kinetics of activation and inactivation in this version of TREK-2 (Fig. S2). However, for TWIK-1*, although the rate of channel activation also decreased at 3°C , the inactivation kinetics of TWIK-1 were still unable to be resolved (Fig. 3D and Fig. S2).

The SF is highly unstable in the presence of K⁺

Based on these observations, we hypothesized that when the SF is occupied by K⁺, it may be extremely unstable, resulting in almost instantaneous inactivation when Rb⁺ is replaced by K⁺. To further investigate this, Rb⁺ was used as the permeant ion, and increasing concentrations of K⁺ were added. Intriguingly, we found that very low concentrations of K⁺ strongly inhibited these Rb⁺ currents (Fig. 4A). A full dose–response curve for this effect revealed an IC₅₀ for K⁺ of 2.8 mM (Fig. 4A). In marked contrast, the TREK-1 K2P channel showed comparatively weaker K⁺-dependent inhibition of Rb⁺ currents (IC₅₀ of ~ 68 mM). To test whether this observed inhibitory effect of K⁺ on WT* TWIK-1 currents is specific, we compared the effect of several other monovalent cations. We found that K⁺ exhibited the strongest inhibitory effect and Cs⁺ the weakest (Fig. 4B).

Unusual filter gating properties of TWIK-1




Figure 4. High-affinity inhibition of TWIK-1* Rb⁺ currents by intracellular K⁺. *A*, comparison of K⁺ inhibition of Rb⁺ currents in TWIK-1* and TREK-1. *Top panels*, in both cases, the intracellular concentration of Rb⁺ is kept constant at 120 mM, and increasing concentrations of K⁺ are added. Currents were recorded in response to a voltage ramp from -80 to +100 mV. In all cases, these currents can be blocked by intracellular application of 1 mM TPA (example shown in red). *Bottom panels*, TWIK-1* and TREK-1 currents from the experiments in the *top panels* are plotted against intracellular K⁺ concentrations. *B*, comparison of the inhibitory effect of different monovalent cations on TWIK-1* Rb⁺ currents. Currents were elicited in response to voltage ramps from -80 to +120 mV. K⁺ is the most potent inhibitory ion within the physiological voltage range. Note that Cs⁺, which usually acts as a permeant blocker of most K⁺ channels, has the least inhibitory effect.

This also appears to be in agreement with the relative activatory effect of these different ions shown in Fig. 3A.

Contribution of the selectivity filter gate

We next set out to determine the mechanisms underlying this K⁺-dependent block of Rb⁺ permeation. In several previous studies, we have shown that most forms of gating in K₂P channels occur primarily within the SF (21, 23–25, 27). Interestingly, the SF of TWIK-1 exhibits several differences compared with other K₂P channels within the canonical filter sequence as well as a dynamic K⁺ > Na⁺ selectivity that can be modulated by mutations close to the filter (15). In that study, it was reported that the T118I mutation in the




Figure 5. Inhibitory effect of extracellular K⁺ on whole-cell Rb⁺ currents. *A*, inhibition of WT TWIK-1* by increasing concentrations of extracellular K⁺. Channels were expressed in HEK293 cells, and the extracellular Rb⁺ concentration was kept constant (150 mM) with increasing concentrations of K⁺ as indicated. *Right panel*, an example dose–response curve determined from the adjacent traces. *B*, a structural cartoon depicting the two pore loops of TWIK-1 and the relative position of the mutations with the SF. *C*, scatterplots of individual IC₅₀ values determined from independent experiments similar to those shown in *A*. Data are shown for WT TWIK-1* and for the T118I, L108F/F109Y, and L228F mutants, which stabilize the dynamic K⁺/Na⁺ selectivity of TWIK-1 (15). Similar data are also shown for mutation of the two conserved threonines in the canonical K⁺-selective filter sequences of the first and second P-loops. The *errors bars* represent the mean and standard deviation. *D*, the difference in the time course of K⁺ block and release from K⁺ block is consistent with tight binding of K⁺ within the pore. Representative traces show the rate of K⁺ block (black) and release from block (blue) as K⁺ or Rb⁺ solutions are applied, respectively. *Right panel*, the values recorded from multiple individual experiments (see also Fig. S3).

first P-loop (T₁GYG) “stabilized” the channel in its K⁺-selective conformation, whereas the L108F/F109Y mutation in the preceding pore helix stabilized its Na⁺-selective conformation.

We therefore examined the effect of these two mutations on K⁺-dependent inhibition of WT TWIK-1* Rb⁺ currents. To perform these experiments, we expressed TWIK-1* in mammalian HEK293 cells and measured inward currents in the whole-cell configuration. Because Rb⁺ also activates the SF gate from the extracellular side (Fig. 2C) we measured the inhibitory effect of extracellular K⁺ on these inwardly directed Rb⁺ currents. Fig. 5A shows that, in this recording configuration, WT TWIK-1* also still exhibited clear K⁺-dependent

inhibition of these inward Rb^+ currents, albeit with reduced affinity (IC_{50} , ~ 22 mM).

To examine the role of the SF, we studied the effect of K^+ block on several mutants within the SF of TWIK-1* (Fig. 5B). The T118I mutation in TWIK-1* has been shown previously to stabilize the filter in an apparently K^+ -selective conformation (Fig. 5B). Consistent with this, we found that this mutant exhibited the same level of K^+ -dependent block as WT TWIK-1*, as did the L228F mutation, which restores the consensus sequence of the second P-loop (Fig. 5C) and also does not affect TWIK-1* K^+ selectivity (15). However, the L108F/F109Y mutation within pore helix 1 (Fig. 5B), which has been reported to reduce K^+ selectivity (15), markedly reduced this inhibitory effect (IC_{50} , ~ 62 mM) (Fig. 5C).

Functional asymmetry within the filter

Overall, these results strongly suggest a direct interaction of K^+ with the SF of TWIK-1 that induces an unstable and therefore poorly conductive conformation of the SF gate. Interestingly, a recent study suggested that K^+ exhibits a relatively low affinity for TWIK-1 (28). However, TWIK-1 activity was not measured directly, and ion exchange-induced difference IR spectroscopy of purified TWIK-1 channels in the absence of any transmembrane electrochemical potential was used to determine this interaction. In contrast, our approach provides a more direct measurement of how different ions permeate through the SF. Nevertheless, to further support our observations, we examined the role of the filter gate in greater detail.

In a previous study of K2P channel gating, we have shown that, like KcsA, mutations within the conserved threonines of the consensus TXGYG sequence can have a profound effect on the SF gating mechanism (23). These conserved threonines are located at the lower entrance to the filter, with their hydroxyl groups partially substituting for the waters of hydration of the permeant ion (usually K^+) immediately prior to the ion entering the SF. We therefore mutated these two threonine residues and examined how they influence the high-affinity K^+ block of Rb^+ currents through TWIK-1* channels.

We found that mutation of the first conserved threonine (Thr¹¹⁷) had no effect (Fig. 5C), whereas mutation of Thr²²⁵ in the second P-loop markedly reduced this K^+ -dependent inhibition. This result not only supports the role of the SF in this interaction of K^+ with TWIK-1 but also highlights the functionally asymmetric contribution of these two pore-forming loops to the SF gate.

To further support this inhibitory effect of K^+ , we also examined the kinetics of inhibition on WT TWIK-1* channels. Although the absolute values determined depend on the limitations of the perfusion system, we found that the relative rates of blocking and unblocking were different. Consistent with a tight binding of K^+ within the pore, the rate of K^+ inhibition was faster (τ , ~ 450 ms) than recovery of these currents upon washout (τ , ~ 1 s) (Fig. 5D and Fig. S3).

Differential occupancy of K^+ and Rb^+ in the SF

To gain further insight into the mechanism by which K^+ and Rb^+ may interact differently within the filter of TWIK-1 to influence the SF gate, we employed atomistic double-bilayer




Figure 6. MD simulation reveals differential occupancy of SF by K^+ versus Rb^+ . Double-bilayer simulations of TWIK-1 with either K^+ or Rb^+ reveal a different occupancy of the S1–S4 ion-binding sites. The relative ion occupancy for outward currents is shown plotted against the position of the SF on the z axis (K^+ in black and Rb^+ in red). The data shown were collected from multiple repeats that each represent a total of 6- μs simulation during which seven outward permeation events were observed for K^+ and 13 events for Rb^+ . The peaks shown on the left correspond to the S1–S4 ion-binding sites. These are structurally aligned with a depiction of the selectivity filter and pore helices of TWIK-1.

molecular dynamics (MD) simulations of permeation similar to those we used recently to examine permeation in other types of K^+ channels, including K2P channels (23, 24, 29). Simulations with either K^+ or Rb^+ as the permeant ion were conducted (Fig. 6). Over a total 6- μs simulation of outward permeation, the four main ion binding sites (S1–S4) within the filter of TWIK-1 displayed a markedly different occupancy by K^+ compared with Rb^+ . In particular, although K^+ exhibited preferential occupancy of S2 and S4 and relatively little occupancy at S1 and S3, Rb^+ occupied all four sites, including S1 and S3. These results support previous studies that showed that K^+ and Rb^+ interact differently with the SF in other K^+ channels (23, 30) and suggest that a differential interaction of permeant ions at the S1 and S3 sites within TWIK-1 may contribute to its unusual functional properties.

Discussion

Overall, our results suggest that, when present in the plasma membrane under normal physiological conditions and voltages, K^+ stabilizes the TWIK-1 SF gate in an unstable, poorly conductive conformation. This is also supported by most functional studies that have attempted to record the functional activity of TWIK-1 using K^+ as the permeant ion. Nevertheless, several previous studies have shown that the levels of whole-cell TWIK-1 channel K^+ current can be enhanced by removal of an endocytic retrieval motif within the C terminus (*i.e.* the TWIK-1* mutant used in this study), and our results are consistent with this observation, although these effects appear to be via changes in cell surface expression rather than altering the gating properties of the channel *per se*. Other studies have also shown modest increases in whole-cell K^+ currents of both WT TWIK-1 and TWIK-1* when Lys²⁷⁴ within the C-helix is mutated. Again, our results are also consistent with this observation, although the underlying mechanism still remains controversial. Some studies have proposed that posttranslational sumoylation of Lys²⁷⁴ somehow silences channel activity, whereas other studies suggest that sumoylation is not respon-

Unusual filter gating properties of TWIK-1

sible. Our results show that mutation of other charged residues within the C-helix can also produce similar increases in TWIK-1* whole-cell current, suggesting that the effect of mutating Lys²⁷⁴ may not be as unique as thought previously. Either way, the increases in current seen with these C-helix mutations are relatively small compared with mutations within the hydrophobic cuff of the inner cavity (e.g. L146N). However, even the effect of the L146N mutation is small compared with the large and robust currents that can be obtained when Rb⁺ is used as the permeant ion. This suggests that the unusual properties of the SF gate dominate the low intrinsic open probability of TWIK-1 when the channel is present in the plasma membrane.

In support of this, we show that, like other K2Ps, TWIK-1 does indeed possess a flux-gating mechanism but that it handles K⁺ and Rb⁺ differently and exhibits significantly altered properties compared with most other K2P channels. In particular, it displays a different ion activation profile, and with K⁺ this voltage-dependence is shifted into the supra-physiological range. Another major difference is that the SF gate can be activated by extracellular Rb⁺ and exhibits a highly unusual K⁺-dependent block. Our results demonstrate that these unusual properties are conferred by the SF and that the contributions of the two pore-forming loops are unequal. This may be the result of an asymmetric contribution of the two P-loops to K2P channel gating and/or the unusual sequence of the consensus K⁺ filter motifs found in TWIK-1.

The physiological significance of these biophysical differences is unclear. One possibility may be related to the dynamic nature of TWIK-1 ion selectivity and that significant levels of TWIK-1 channels have been reported within endosomes, where the unusual ionic environment and pH may modulate its functional properties (15, 18, 31). Also, the K⁺ concentrations within intracellular endosomes have been reported to be highly dynamic and proposed to play a role in the control of viral trafficking (32). In addition, K2P channels are thought to play a key role in this process, although a specific role for TWIK-1 remains to be determined (33). Nevertheless, the unique biophysical properties we have identified in this study will clearly aid future functional and cellular studies of this enigmatic ion channel.

Experimental procedures

Electrophysiological studies of TWIK-1 and TREK-1 were performed as described previously (12, 15). Briefly, human TWIK-1 was subcloned into pFAW for expression oocytes and HEK293 cells. Two-electrode voltage clamp recordings of whole-cell currents in *Xenopus* oocytes were recorded in ND96 solution at pH 7.4 (96 mM NaCl, 2 mM KCl, 2 mM MgCl₂, 1.8 mM CaCl₂, and 5 mM HEPES). For excised patch recordings, the intracellular solution (K⁺_{int}) had the following composition: 120 mM KCl, 10 mM HEPES, and 2 mM EGTA, adjusted to the appropriate pH with HCl/KOH. The extracellular solution (K⁺_{ex}, pipette solution) comprised 120 mM KCl, 10 mM HEPES, and 3.6 mM CaCl₂ adjusted to pH 7.4 by HCl/KOH. In relevant recordings, the extra- and intracellular K⁺ was replaced by Cs⁺, Na⁺, NH₄⁺, Rb⁺, or Tl⁺, and the pH was adjusted by hydroxide of the relevant ion species and HCl. For recordings in HEK293

cells, solutions had the following composition: 150 mM KCl and 10 mM HEPES for extracellular solutions, and the pH was adjusted to 7.4 by HCl/KOH. Rb⁺ replaced K⁺ in other solutions and pH was adjusted by RbOH/HCl. Intracellular solutions were composed of 140 mM KCl, 10 mM HEPES, and 2 mM EGTA. The pH was adjusted to 8.0. All solution exchanges were done using a gravity-driven multibarrel micropipette.

For expression in oocytes, 50 nl of mRNA (500 ng/μl) was manually injected into Dumont stage VI oocytes and incubated at 17 °C for 1–3 days prior to use. HEK293 cells were cultivated in DMEM and Ham's F-12 nutrient mixture and seeded in 5% gelatin-treated Petri dishes 24 h prior to transfection. The transfection was done using FuGENE[®] HD transfection reagent according to the manufacturer's protocol. Recordings were made 24–48 h after transfection.

For excised patches from oocytes, pipettes were made using thick-walled borosilicate glass with a resistance of 0.3–0.9 megaohm and filled with the extracellular solutions. HEK293 cell experiments were done in whole-cell configuration using extra-thin glass pipettes with a resistance of 1–1.9 megaohm and filled with intracellular solutions. All pipettes were purchased from Science Products GmbH. Currents were recorded with an EPC10 amplifier running under Patchmaster software (HEKA Electronics, Lamprecht, Germany) for excised patches or Axopatch[™] 200B running under Clampex (Molecular Devices) and sampled at 10 kHz and 250 kHz, respectively, with the analog filter set to 3 kHz and 10 kHz, respectively, and stored for later analysis. IC₅₀ values were calculated by fitting data with the Hill equation: $\text{base} + (\text{max} - \text{base})/[1 + (\text{xhalf}/\text{x})^h]$, where base is the zero current, max is the maximum current, x is the ligand concentration, xhalf is the concentration for half-maximal occupation of the ligand-binding site (inhibition by TPA or H⁺), and h is the Hill coefficient. The voltage-activated traces were fitted by the exponential function $y_0 + A \exp[-(t - t_0)/\tau]$, where y₀ is the current where the fitting starts, A, the fit constant, t₀ the time where the fitting starts, and τ the time constant for the current increase. The time constant for block or release from the block are calculated by fitting the traces with the sigmoidal function $F(t) = A_{\text{base}} + \{(A_{\text{max}} - A_{\text{base}})/[1 + (t_{1/2}/t)^n]\}$, where A_{base} is the base current, A_{max} is the maximum current, t is the time, and t_{1/2} is the time where the current reaches half the maximum current (A_{max}), and n is the rise rate. Examples of the fits used are shown in the supplemental information (Fig. S3). Unless otherwise described, values result from at least three independent experiments given as mean ± S.E.

MD simulations were performed as described before (23, 24, 29, 34). Briefly, we used a modified version of GROMACS 4.6, the AMBER99sb force field, and the SPC/E water model for the equilibrium and production simulations. The simulations were performed at 300 K with a velocity-rescaling thermostat, and the pressure was kept at 1 bar using a semi-isotropic Berendsen barostat. All bonds were constrained with the LINCS algorithm, and using virtual sites for hydrogen atoms allowed simulations to be performed with a 4-fs integration time step. The crystallographic structure of TWIK-1 (PDB code 3UKM) (9) was adopted as the initial structure in the MD simulations and was embedded into an aqueous 1-palmitoyl-2-oleoyl-sn-

glycero-3-phosphocholine (POPC) lipid bilayer with 0.6 M KCl. The system was equilibrated for 20 ns with position restraints on all heavy atoms using a force constant of $1000 \text{ kJ mol}^{-1} \text{ nm}^{-2}$ to the reference structure, followed by an additional 10-ns simulation without position restraints. Production runs were then performed on the equilibrated TWIK-1 structure using a computational electrophysiology protocol (34). For K^+ and Rb^+ , 20 replicas of 300 ns were carried out for each ion type. Potential gradients were generated by introducing a charge difference of two K^+ or two Rb^+ ions between the two compartments separated by the two lipid bilayers. During the MD simulations, the number of the ions was kept constant by an additional algorithm (34). The potential difference between two compartments corresponds to $360 \pm 70 \text{ mV}$ and $390 \pm 70 \text{ mV}$ for K^+ and Rb^+ simulations, respectively.

Author contributions—E. N.-A. data curation; E. N.-A., H. S., M. S., and S. J. T. formal analysis; E. N.-A., F. A.-W., F. C. C., H. S., and M. S. investigation; E. N.-A., F. A.-W., F. C. C., and T. B. methodology; E. N.-A. and S. J. T. writing-original draft; E. N.-A., F. A.-W., F. C. C., H. S., M. S., T. B., and S. J. T. writing-review and editing; H. S. and S. J. T. visualization; T. B. and S. J. T. conceptualization; T. B. and S. J. T. resources; T. B. and S. J. T. supervision; T. B. and S. J. T. funding acquisition; T. B. and S. J. T. project administration; S. J. T. validation.

References

- Enyedi, P., and Czirják, G. (2010) Molecular background of leak K^+ currents: two-pore domain potassium channels. *Physiol. Rev.* **90**, 559–605 [CrossRef Medline](#)
- Niemeyer, M. I., Cid, L. P., González, W., and Sepúlveda, F. V. (2016) Gating, regulation, and structure in K2P K^+ channels: *in varietate concordia*? *Mol. Pharmacol.* **90**, 309–317 [CrossRef Medline](#)
- Gada, K., and Plant, L. D. (2019) Two-pore domain potassium channels: emerging targets for novel analgesic drugs: IUPHAR Review 26. *Br. J. Pharmacol.* **176**, 256–266 [CrossRef Medline](#)
- Lesage, F., Guillemare, E., Fink, M., Duprat, F., Lazdunski, M., Romey, G., and Barhanin, J. (1996) TWIK-1, a ubiquitous human weakly inward rectifying K^+ channel with a novel structure. *EMBO J.* **15**, 1004–1011 [CrossRef Medline](#)
- Goldstein, S. A. (2011) K2P potassium channels, mysterious and paradoxically exciting. *Sci. Signal.* **4**, pe35 [Medline](#)
- Yarishkin, O., Lee, D. Y., Kim, E., Cho, C. H., Choi, J. H., Lee, C. J., Hwang, E. M., and Park, J. Y. (2014) TWIK-1 contributes to the intrinsic excitability of dentate granule cells in mouse hippocampus. *Mol. Brain* **7**, 80 [CrossRef Medline](#)
- Ma, L., Zhang, X., and Chen, H. (2011) TWIK-1 two-pore domain potassium channels change ion selectivity and conduct inward leak sodium currents in hypokalemia. *Sci. Signal.* **4**, ra37 [CrossRef Medline](#)
- Rajan, S., Plant, L. D., Rabin, M. L., Butler, M. H., and Goldstein, S. A. (2005) Sumoylation silences the plasma membrane leak K^+ channel K2P1. *Cell* **121**, 37–47 [CrossRef Medline](#)
- Miller, A. N., and Long, S. B. (2012) Crystal structure of the human two-pore domain potassium channel K2P1. *Science* **335**, 432–436 [CrossRef Medline](#)
- Feliciangeli, S., Bendahhou, S., Sandoz, G., Gounon, P., Reichold, M., Warth, R., Lazdunski, M., Barhanin, J., and Lesage, F. (2007) Does sumoylation control K2P1/TWIK1 background K^+ channels? *Cell* **130**, 563–569 [CrossRef Medline](#)
- Feliciangeli, S., Tardy, M. P., Sandoz, G., Chatelain, F. C., Warth, R., Barhanin, J., Bendahhou, S., and Lesage, F. (2010) Potassium channel silencing by constitutive endocytosis and intracellular sequestration. *J. Biol. Chem.* **285**, 4798–4805 [CrossRef Medline](#)
- Aryal, P., Abd-Wahab, F., Bucci, G., Sansom, M. S., and Tucker, S. J. (2014) A hydrophobic barrier deep within the inner pore of the TWIK-1 K2P potassium channel. *Nat. Commun.* **5**, 4377 [CrossRef Medline](#)
- Aryal, P., Abd-Wahab, F., Bucci, G., Sansom, M. S., and Tucker, S. J. (2015) Influence of lipids on the hydrophobic barrier within the pore of the TWIK-1 K2P channel. *Channels* **9**, 44–49 [CrossRef Medline](#)
- Aryal, P., Sansom, M. S., and Tucker, S. J. (2015) Hydrophobic gating in ion channels. *J. Mol. Biol.* **427**, 121–130 [CrossRef Medline](#)
- Chatelain, F. C., Bichet, D., Douguet, D., Feliciangeli, S., Bendahhou, S., Reichold, M., Warth, R., Barhanin, J., and Lesage, F. (2012) TWIK1, a unique background channel with variable ion selectivity. *Proc. Natl. Acad. Sci. U.S.A.* **109**, 5499–5504 [CrossRef Medline](#)
- Ben Soussia, I., El Mouridi, S., Kang, D., Leclercq-Blondel, A., Khoubza, L., Tardy, P., Zariohi, N., Gendrel, M., Lesage, F., Kim, E. J., Bichet, D., Andriani, O., and Boulin, T. (2019) Mutation of a single residue promotes gating of vertebrate and invertebrate two-pore domain potassium channels. *Nat. Commun.* **10**, 787 [CrossRef Medline](#)
- Ma, L., Xie, Y. P., Zhou, M., and Chen, H. (2012) Silent TWIK-1 potassium channels conduct monovalent cation currents. *Biophys. J.* **102**, L34–L36 [CrossRef Medline](#)
- Chen, H., Chatelain, F. C., and Lesage, F. (2014) Altered and dynamic ion selectivity of K^+ channels in cell development and excitability. *Trends Pharmacol. Sci.* **35**, 461–469 [CrossRef Medline](#)
- Bagriantsev, S. N., Peyronnet, R., Clark, K. A., Honoré, E., and Minor, D. L., Jr. (2011) Multiple modalities converge on a common gate to control K2P channel function. *EMBO J.* **30**, 3594–3606 [CrossRef Medline](#)
- Ben-Abu, Y., Zhou, Y., Zilberberg, N., and Yifrach, O. (2009) Inverse coupling in leak and voltage-activated K^+ channel gates underlies distinct roles in electrical signaling. *Nat. Struct. Mol. Biol.* **16**, 71–79 [CrossRef Medline](#)
- Piechotta, P. L., Rapedius, M., Stansfeld, P. J., Bollepalli, M. K., Ehrlich, G., Ehrlich, G., Andres-Enguix, I., Fritzenschaft, H., Decher, N., Sansom, M. S., Tucker, S. J., and Baukrowitz, T. (2011) The pore structure and gating mechanism of K2P channels. *EMBO J.* **30**, 3607–3619 [CrossRef Medline](#)
- Zilberberg, N., Ilan, N., and Goldstein, S. A. (2001) KCNKO: opening and closing the 2-P-domain potassium leak channel entails “C-type” gating of the outer pore. *Neuron* **32**, 635–648 [CrossRef Medline](#)
- Schewe, M., Nematian-Ardestani, E., Sun, H., Musinszki, M., Cordeiro, S., Bucci, G., de Groot, B. L., Tucker, S. J., Rapedius, M., and Baukrowitz, T. (2016) A non-canonical voltage-sensing mechanism controls gating in K2P K^+ channels. *Cell* **164**, 937–949 [CrossRef Medline](#)
- Schewe, M., Sun, H., Mert, Ü., Mackenzie, A., Pike, A. C. W., Schulz, F., Constantin, C., Vowinkel, K. S., Conrad, L. J., Kiper, A. K., Gonzalez, W., Musinszki, M., Tegtmeyer, M., Pryde, D. C., Belabed, H., et al. (2019) A pharmacological master key mechanism that unlocks the selectivity filter gate in K^+ channels. *Science* **363**, 875–880 [CrossRef Medline](#)
- Rapedius, M., Schmidt, M. R., Sharma, C., Stansfeld, P. J., Sansom, M. S., Baukrowitz, T., and Tucker, S. J. (2012) State-independent intracellular access of quaternary ammonium blockers to the pore of TREK-1. *Channels* **6**, 473–478 [CrossRef Medline](#)
- Dong, Y. Y., Pike, A. C., Mackenzie, A., McClenaghan, C., Aryal, P., Dong, L., Quigley, A., Grieben, M., Goubin, S., Mukhopadhyay, S., Ruda, G. F., Clausen, M. V., Cao, L., Brennan, P. E., Burgess-Brown, N. A., et al. (2015) K2P channel gating mechanisms revealed by structures of TREK-2 and a complex with Prozac. *Science* **347**, 1256–1259 [CrossRef Medline](#)
- Nematian-Ardestani, E., Jarerattanachai, V., Aryal, P., Sansom, M. S. P., and Tucker, S. J. (2017) The effects of stretch activation on ionic selectivity of the TREK-2 K2P K^+ channel. *Channels* **11**, 482–486 [CrossRef Medline](#)
- Tsukamoto, H., Higashi, M., Motoki, H., Watanabe, H., Ganser, C., Nakajo, K., Kubo, Y., Uchihashi, T., and Furutani, Y. (2018) Structural properties determining low K^+ affinity of the selectivity filter in the TWIK1 K^+ channel. *J. Biol. Chem.* **293**, 6969–6984 [CrossRef Medline](#)
- Köpfer, D. A., Song, C., Gruene, T., Sheldrick, G. M., Zachariae, U., and de Groot, B. L. (2014) Ion permeation in K^+ channels occurs by direct Coulomb knock-on. *Science* **346**, 352–355 [CrossRef Medline](#)
- Zhou, Y., and MacKinnon, R. (2003) The occupancy of ions in the K^+ selectivity filter: charge balance and coupling of ion binding to a protein

Unusual filter gating properties of TWIK-1

- conformational change underlie high conduction rates. *J. Mol. Biol.* **333**, 965–975 [CrossRef Medline](#)
31. Bobak, N., Feliciangeli, S., Chen, C. C., Ben Soussia, I., Bittner, S., Pagnotta, S., Ruck, T., Biel, M., Wahl-Schott, C., Grimm, C., Meuth, S. G., and Lesage, F. (2017) Recombinant tandem of pore-domains in a weakly inward rectifying K⁺ channel 2 (TWIK2) forms active lysosomal channels. *Sci. Rep.* **7**, 649 [CrossRef Medline](#)
 32. Hover, S., Foster, B., Fontana, J., Kohl, A., Goldstein, S. A. N., Barr, J. N., and Mankouri, J. (2018) Bunyavirus requirement for endosomal K⁺ reveals new roles of cellular ion channels during infection. *PLoS Pathog.* **14**, e1006845 [CrossRef Medline](#)
 33. Hover, S., King, B., Hall, B., Loundras, E. A., Taqi, H., Daly, J., Dallas, M., Peers, C., Schnettler, E., McKimmie, C., Kohl, A., Barr, J. N., and Mankouri, J. (2016) Modulation of potassium channels inhibits bunyavirus infection. *J. Biol. Chem.* **291**, 3411–3422 [CrossRef Medline](#)
 34. Kutzner, C., Grubmüller, H., de Groot, B. L., and Zachariae, U. (2011) Computational electrophysiology: the molecular dynamics of ion channel permeation and selectivity in atomistic detail. *Biophys. J.* **101**, 809–817 [CrossRef Medline](#)

**Selectivity filter instability dominates the low intrinsic activity of the TWIK-1 K²P
K⁺ channel**

Ehsan Nematian-Ardestani, Firdaus Abd-Wahab, Franck C. Chatelain, Han Sun, Marcus Schewe, Thomas Baukowitz and Stephen J. Tucker

J. Biol. Chem. 2020, 295:610-618.

doi: 10.1074/jbc.RA119.010612 originally published online December 5, 2019

Access the most updated version of this article at doi: [10.1074/jbc.RA119.010612](https://doi.org/10.1074/jbc.RA119.010612)

Alerts:

- [When this article is cited](#)
- [When a correction for this article is posted](#)

[Click here](#) to choose from all of JBC's e-mail alerts

This article cites 34 references, 11 of which can be accessed free at <http://www.jbc.org/content/295/2/610.full.html#ref-list-1>

Supporting Information


Selectivity filter instability dominates the low intrinsic activity of the TWIK-1 K2P K⁺ channel

Ehsan Nematian-Ardestani¹, M. Firdaus Abd-Wahab¹, Franck C. Chatelain², Han Sun³, Marcus Schewe⁴, Thomas Baukrowitz⁴ and Stephen J Tucker^{1*}


From the ¹Clarendon Laboratory, Department of Physics, University of Oxford, Oxford OX1 3PU, United Kingdom; ²Université Côte d'Azur, Centre National de la Recherche Scientifique, Institut de Pharmacologie Moléculaire et Cellulaire, Labex ICST, Valbonne, France; ³Leibniz-Forschungsinstitut für Molekulare Pharmakologie (FMP), Department of Structural Biology, 13125 Berlin, Germany, and ⁴Institute of Physiology, Christian-Albrechts University of Kiel, 24118 Kiel, Germany.

Running Title: Unusual filter gating properties of TWIK-1


*Corresponding Author: Stephen J. Tucker, Clarendon Laboratory, Department of Physics, University of Oxford, OX1 3PU, United Kingdom. E-mail: stephen.tucker@physics.ox.ac.uk



Supplemental Fig. S1. Example currents for WT TWIK-1* and the three key mutations shown in Figure 1B. These mutants all produce increases in whole-cell current that are equivalent to the effect of the previously reported K274E mutation (8). Currents were recorded by TEVC in response to voltage steps from -120 to +60 mV from a holding potential of -80 mV. Reported values were measured at 0 mV.



Supplemental Fig. S2. A) Example of the fits used (red) on Rb^+ activated TWIK-1 currents. These fits were used to obtain the rate constants shown in Fig. 3C which demonstrate that activation is not voltage-dependent (see methods for details). B) Currents recorded in excised patches from a mutant version of TREK-2 (Gly⁶⁷-Glu³⁴⁰) (26) at both 23° and 3° under the same conditions shown in Fig. 3D. At 23° no tail currents are visible, but upon cooling to 3° the rate of activation slows, as well as the tail currents so they become more visible. An overlay of these currents at +100 mV is shown in the right hand panel.



Supplemental Fig. S3 Example of the fits used (in red) to obtain the rate constants for K^+ block and unblock that are shown in Fig. 5D. See experimental procedures for details.